

Critical temperature and phase diagram for a 2D superfluid Fermi-gas with repulsion.

D.V. Efremov^a, M.S. Mar'enko^{a,1}, M.A. Baranov^b, M.Yu. Kagan^a

^a*P.L.Kapitza Institute for Physical Problems, Moscow 117334, Russia*

^b*Russian Research Center "Kurchatov Institute", Moscow 123182, Russia*

Abstract

In the framework of a 2D Fermi-gas with short-range repulsive interaction we find the critical temperature of the superfluid phase transition based on Kohn-Luttinger effect, and analyze its dependence on magnetic field. We calculate strong-coupling corrections to the Ginzburg-Landau free energy functional and analyze the stability of 2D superfluid phases. Possible applications of the results to real systems are discussed.

Keywords: Two-dimensional Fermi-liquid, ³He-⁴He mixtures, superfluidity, superconductivity.

1. Introduction

Recent experiments on ³He submonolayers absorbed on graphite [1], ³He atoms bound to the surface of ⁴He films [2–4] and discovery of superconductivity in layered HTSC materials stimulate theoretical studies of superfluid phase transition in a 2D Fermi liquid. It was shown [5,6], that even in the case of a bare repulsive *s*-wave interaction in a 3D Fermi system, there is a transition to the *p*-wave superfluid state due to a singularity in the effective interaction near the momentum transfer $q = 2p_F$ (Kohn-Luttinger effect), $\Gamma_{\text{sing}}(q) \sim (2p_F - q) \ln |2p_F - q|$, where p_F is the Fermi momentum. Further analysis [7] shows that the similar effect takes place also in 2D case.

2. Theoretical model and results

We consider a 2D non-ideal Fermi gas with a short-range repulsive interaction, $r_0 p_F \ll 1$, where r_0 is the range of potential, such that the perturbative expansion in a gas parameter f_0 , related to the *s*-wave scattering amplitude between two particles on the Fermi surface, is legitimate. For the considered case the gas parameter is given by

$$f_0 = [4\pi/mU_0 + \ln(r_0 p_F)^{-2}]^{-1},$$

where U_0 is the *s*-component of the potential.

It turns out that for a 2D Fermi gas the Kohn-Luttinger mechanism works in the third order in f_0 , because the singular contribution of the second order in f_0 to the effective interaction,

$$(m/4\pi)\Gamma_{\text{sing}}^{(2)}(q) = -f_0^2 \text{Re} \sqrt{1 - (2p_F/q)^2},$$

¹ Corresponding author. Present address: P.L.Kapitza Institute for Physical Problems, Kosygin str. 2, Moscow 117334, Russia. E-mail: maxim@kapitza.ras.ru




Fig. 1. The p -wave critical temperature $T_{c1}/\tilde{\epsilon}$ as a function of polarization $\alpha = \frac{(n_{\uparrow} - n_{\downarrow})}{(n_{\uparrow} + n_{\downarrow})}$ for typical $f_0 = 0.3$.

is zero for $q \leq 2p_F$, and, therefore, does not contribute to the p -harmonic. The result for $\Gamma_{\text{sing}}(q)$ in the third order in f_0 is

$$(m/4\pi)\Gamma_{\text{sing}}^{(3)}(q) \sim -f_0^3 \text{Re} \sqrt{1 - (q/2p_F)^2},$$

and it leads to an effective attraction in the p -wave channel. Numerical calculations of third-order diagrams give for the p -wave critical temperature

$$T_{c1} = \tilde{\epsilon} \exp\left(-\frac{1}{6.1f_0^3}\right), \quad (1)$$

where $\tilde{\epsilon} = A \cdot \epsilon_F$, A is the unknown numerical prefactor, $\epsilon_F = p_F^2/2m$ is the Fermi energy.

The typical value of scattering amplitude obtained from the experiments on ^3He atoms on the surface of ^4He is $f_0 \leq 0.3$ for ^3He coverages corresponding to the dilute Fermi gas situation. In this regime the critical temperature is $T_{c1} \sim 10^{-3} K$.

The critical temperature T_{c1} can be significantly increased by applying an external magnetic field, because in this case the effective interaction contributes to T_{c1} already in the second order. The separation of polarization and pairing effects leads to a strongly non-monotonic dependence of the critical temperature on the polarization (Fig. 1).

In the weak coupling (BCS) limit for zero magnetic field, two different phases correspond to the minimum of the free energy (the axial and the planar phases) [8]. To lift this degeneracy one has to consider next order ($\sim T_c/\epsilon_F$) corrections. Following [9], we have calculated the difference between free energies of axial and planar phases to this order:

$$\Phi_{\text{axial}} - \Phi_{\text{planar}} \sim -\frac{N(0)}{T_c^2} \left(\frac{T_c}{\epsilon_F}\right) \times \int \frac{d\varphi}{2\pi} |\sin \varphi| T_a^2(\hat{\mathbf{k}}_1, \hat{\mathbf{k}}_2; \hat{\mathbf{k}}_1, \hat{\mathbf{k}}_2), \quad (2)$$

where $N(0)$ is the density of states on the Fermi surface, $T_a(\hat{\mathbf{k}}_1, \hat{\mathbf{k}}_2; \hat{\mathbf{k}}_3, \hat{\mathbf{k}}_4)$ the spin-antisymmetric part of quasiparticle scattering amplitude with momenta $\mathbf{k}_i$ on the Fermi surface and φ the angle between $\hat{\mathbf{k}}_1$ and $\hat{\mathbf{k}}_2$. Clearly, the energy difference (2) is always negative, therefore in the absence of a magnetic field the superfluid Fermi-gas forms the axial phase.

Acknowledgement

The authors are grateful to A.V. Chubukov, G. Frossati, H. Godfrin, R.B. Hallock, J. Saunders, J. Nyeki and B.P. Cowan for stimulating discussions. This work has been supported by RFBR grants 98-02-17077, 97-02-16532 and INTAS grant 98-963.

References

- [1] D.S. Greywall, Phys. Rev B **41** (1990) 1842.
- [2] M. Dann, J. Nyeki, B.P. Cowan, J. Saunders, Phys. Rev Lett. **82** (1999) 4030.
- [3] D.T. Sprague, N. Alikasem, P.A. Sheldon, R.B. Hallock, Phys. Rev Lett. **72** (1994) 384.
- [4] K.-D. Morhard, C. Bäuerle, J. Bossy, Yu.M. Bunkov, S.N. Fisher, H. Godfrin, Phys. Rev. B **53** (1996) 2658.
- [5] W. Kohn, J.H. Luttinger, Phys. Rev Lett. **15** (1965) 524.
- [6] M.Yu. Kagan, A.V. Chubukov, JETP Lett. **47** (1988) 525.
- [7] M.A. Baranov, A.V. Chubukov, M.Yu. Kagan, Int. J. Mod. Phys. B **6** (1992) 2471, A.V. Chubukov, Phys. Rev B **48** (1993) 1097.
- [8] P.N. Brusov, V.N. Popov, JETP **80** (1981) 1564.
- [9] D. Rainer, J.W. Serene, Phys. Rev. B **13** (1976) 4745.